

*Integrated Perspectives: A Journal of Interdisciplinary Research*  
*Indexed by Crossref DOI*  
*Inaugural issue (Vol.1, No.1, Jan-March, 2026)*

*Establishing Climate Variability and Change Using Rainfall, Temperature and Wind as proxies; Evidence from Northwestern Nigeria*

Da'u Abba Umar<sup>1</sup>, Jabir Haruna Abdulkareem<sup>2</sup> Salisu Lawan Halliru<sup>3</sup>, Aliyu Muhammad Inuwa<sup>4</sup>, Nafisa Usman Tafida<sup>5</sup>, Ramatu Dahiru<sup>6</sup>, Binta Zakari<sup>7</sup>, Tukur Usman Garo<sup>8</sup>

<sup>1, 4 & 5</sup> Department of Environmental Sciences, Federal University Dutse, Jigawa State Nigeria

<sup>2</sup> Department of Soil Science, Institute for Agricultural Research Samaru, Ahmadu Bello University Zaria Kaduna, Nigeria

<sup>3</sup> Department of Geography, Yusuf Maitama Sule University of Education Kano, Kano State, Nigeria.

<sup>6</sup> Government Senior Secondary School Jigawar Tsada Dutse, Jigawa State, Nigeria

<sup>7</sup> Department of Geography, Sule Lamido University Kafin Hausa, Jigawa State, Nigeria.

<sup>8</sup> Department of Geography, Aliko Dangote University of Science and Technology Wudil, Kano State, Nigeria

Corresponding author; [abbaumar.d@fud.edu.ng](mailto:abbaumar.d@fud.edu.ng); [daumarkukuma@gmail.com](mailto:daumarkukuma@gmail.com)

## **Abstract**

This study attempts to establish climate change and variability through trend analysis using rainfall, temperature, and wind as proxies. This research intends to determine the geographical and earthly trends of the chosen atmospheric parameters. Data analysis was performed applying the Non-parametric Test, Mann-Kendall Trend Test, and Sen's Slope Estimator. The statistics were prepared and examined using version 29 of the statistical package for social sciences (SPSS) software. Furthermore, Content Analysis (CA) was employed to compare the respondents' perceptions of atmosphere deviation and alteration with the climatic data. The results showed the occurrence of challenges linked to disproportionate rainfall connected to heightened climate variability. This was substantiated through the statistically significant spatial variations in rainfall, temperature, and wind speed, although at different scales. Generally, the results displayed a latitudinal and sequential inclination of rainfall, temperature, and wind speed. However, the time-based variability is more obvious than the longitudinal changes. Yet, this does not mean ignoring the spatial variations the fact that it will affect farmers' livelihoods at different proportion to the magnitude of the horizontal deviations within the study area. Though rainfall plays a significant position in global warming studies due to its influence on climate regulation and food production. Yet the significance of wind and temperature in the climate change mechanism cannot be overlooked considering their crucial roles via evaporation and transpiration.

**Key Words;** *Climate Variability, Northwestern, Nigeria, Rainfall, Temperature*

## **1. Introduction**

Climatological studies reported a global raise in temperatures and variations in precipitation and wind systems in the coming decades (Dan'azumi & Ibrahim, 2023; Zhao et al., 2024). Thus, it is expected globally that various consequences of climate change, such as increasing incidence of heavy rains and erratic precipitations, uncertain onset and cessation of rainy periods, and rising air temperatures will be experienced in the different part of the world. These vagaries in some of the climatic elements terrorized the ecosystems, food production, livelihoods, and infrastructure (Tanko, 2013; Zhao et al., 2024). The changes are not consistent across the country; some regions are heavily affected and therefore more susceptible to climate change than others. It is one of the many difficulties encountered when attempting to differentiate between normal short-term fluctuations and indications of climate change (Roy et al., 2022) that local weather can differ considerably daily, seasonally, and annually.

Climate variability, particularly in terms of rainfall, is a significant feature of the Sahelian atmosphere. Over the past 40 years since 1969, there have been marked declines in average yearly precipitation across the area (Ledda et al., 2024; Pareek, Dhankher, & Foyer, 2020). The Intergovernmental Panel on Climate Change reports that the Sahel region experienced a rainfall reduction of 29–49% from 1968 to 1997 compared to the baseline period of 1931–1960 (O'Neill et al., 2017; Pachauri et al., 2014).

Numerous research have been performed on the forecasting of climatic variability; (Ghalehtimouri, Ros, Rambat, & Nasr, 2024; Njuguna, Onyango, Githaiga, Gituru, & Yan, 2020) examine geographical and earthly patterns of rainstorm variations in Kenya (Mwangi, Julich, Patil, McDonald, & Feger, 2016). Trend Analysis of Droughts during Crop Growing Seasons of Nigeria (Dan'azumi & Ibrahim, 2023). Spatial and Temporal Temperature Trends in Nigeria, 1901-2000 (Umar, Ramli, Aris, Jamil, & Tukur, 2019).

“Detection of Rainfall Trend in Northern Nigeria from 1970 to 2012,” (2025, Sasanya, Adesogan, & Ademola). Examination of changes in rainfall amounts across the northern region of Nigeria (Sawa & Adebayo, 2011). Fluctuation of temperature in Zaria, Northern Nigeria (Ndabula et al., 2013).

The significant climatic variables of rainfall and temperature, commonly analyzed as observed in various studies by numerous scholars, have undergone extensive review over time. Other factors, such as the interplay of wind, rain, and temperature, have yet to be investigated. This research aims to conduct a trend analysis of these variables to reveal the presence of climate variability and change in Northern Nigeria.

## **2. Material and Method**

The study is positioned in the north-western region of the land, wrapped in the middle of Latitudes 11°N to 13°N and Longitudes from 80°E to 10.150°E. This area covered 24,742 km<sup>2</sup> in Jigawa State of Nigeria. According to the 2012 data from the National Bureau of Statistics (Dan’azumi & Ibrahim, 2023), it consists of 27 local administrations and has a population of 4,361,002 people. The State shares its boundaries with Kano and Katsina to the west, Bauchi State to the east, and Yobe State to the northeast. According to the National Population Commission (NPC) of Nigeria, 2009 (White, Uljee, & Engelen, 2012), it splits an international boundary to the north with the Zinder region of the Republic of Niger, as showcase in Figure 1.



The year consists of two main seasons. Humid southwesterly breezes from the Atlantic coast of West Africa cause the monsoon season, which generally lasts from May to September and has included October in recent years. Mustapha et al. (2014) report that the estimated annual rainfall varies between 600 mm and 1,000 mm, with a recent average of around 650 mm.

In some regions of the state, rain is customary for roughly five months. For the rest of the year, a dry season called “Harmattan” dominates, driven by dusty winds from the Sahara desert coming in from the northeast. Umar & Bako (2019) report that average monthly temperatures range from 19 to 35°C, with a peak of 42°C throughout the monsoon season. In the dry season of year maximum and minimum varies between 33°C and 13°C respectively. The average monthly temperatures is between 36°C during the day, and 22°C at night (Umar, Umar, & Tukur, 2017). About four (4) distinct seasons are noticeable: the dry and hot (Bazara), the dry and cool (Rani), the dry and warm (Kaka), and the wet and warm (Damina) seasons (Umar, Ramli, Tukur, Jamil, & Zaudi, 2021).

This study utilized three sets of climatic data that were sourced. The initial set comprises noticed statistics gathered from the general stations of KNARDA (Kano State Agricultural and Rural Development Authority). Over the historical period from 1981 to 1991, eleven (11) zones of the meteorological stations were taken into account. The second data set was gathered from the synoptic stations of JARDA (Jigawa State Agricultural and Rural Development Authority) covering the historical period from 1992 to 2020.

The climatic data gathered from KNARDA and JARDA pertained to rainfall and temperature. To supplement the necessary data for this study, a third set of climatic statistics from MERRA-2 (Modern Era Reanalysis for Research and Applications Version 2) was sourced available at <https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/> from NASA (National Aeronautic and Space

Administration) Langley Research Center (LaRC) Prediction of Worldwide Energy Resource (Power).

Figure 1 provides a brief overview of their location. Data on rainfall, temperature, and wind speed were gathered monthly from NASA. Time series statistics were selected grounded in the standard of the statistics gathered from different stations. Nonetheless, only the National Aeronautics and Space Administration (NASA) offered comprehensive data on rainfall, temperature, and wind speed for the study area between 1981 and 2020. Data missing from observations have been completed with the cross-validation approach as per Ioannis (2017) (Ishaku, Umaru, Adebayo, Löwner, & Okhimamhe, 2024).

The analyses of this research were performed using the Non-parametric Test, Mann-Kendall Test with XLSTAT 2018, and Sen's Slope Estimator under Time Series. The Statistical Package for the Social Sciences (SPSS) Version 29 software was utilized to process and analyze the statistics. Content analysis was applied to examine the connection between the climate data and the perception of respondents regarding climate variability and change.

### **3. Results and Discussion**

#### **3.1 Result for Rainfall Analyses**

Table 1 illustrates the outcomes of the statistical analysis conducted on rainfall data from the study area covering the years 1981 to 2020. In Birniwa, the average annual rainfall amounted to 671.6 mm, with recorded values vary from a lowest of 316.8 mm to a highest of 989.5 mm. The rainfall data for Dutse between 1981 and 2020 shows that the average was 800.5 mm, with a minimum of 263.2 mm and a maximum of 1466.7 mm. The results from Kaugama show similar statistics, with a mean of 700.7 mm, a minimum of 260.9 mm, and a maximum of 1348.1 mm.

Kirikasamma had an average of 594.7 mm, with a minimum of 357.2 mm and a maximum of 969.2 mm. According to the Maigatari results, the average annual precipitation for this timeframe was 512.1 mm, ranging from a low of 346.6 mm to a high of 935.0 mm. The Sule-Tankarkar findings indicated that the average annual precipitation was 604.1 mm, ranging from a minimum of 328.9 mm to a maximum of 1168.0 mm.

**Table 1. Mean Rainfall Data**

<b>Statistic</b>	<b>Birniwa</b>	<b>Dutse</b>	<b>Kaugama</b>	<b>Kirikasamma</b>	<b>Maigatari</b>	<b>Sule Tankarkar</b>
<b>Nbr. of observations</b>	40	40	40	40	40	40
<b>Minimum</b>	316.8	263.2	260.9	357.2	346.6	328.9
<b>Maximum</b>	989.5	1466.7	1348.1	969.2	935.0	1168.0
<b>Range</b>	672.60	1203.51	1087.22	612.01	588.440	839.13
<b>Mean</b>	671.6	800.5	700.7	594.7	512.1	604.1
<b>Variance (n-1)</b>	35136.21	73884.61	51764.03	15491.94	14403.67	22280.72
<b>Standard deviation</b>	187.44	271.81	227.51	124.46	120.015	149.26
<b>Variation coefficient</b>	0.29	0.34	0.32	0.21	0.23	0.24
<b>Skewness (Pearson)</b>	-0.164	0.006	0.230	0.850	1.379	1.326
<b>Kurtosis (Pearson)</b>	-1.01	-0.52	0.14	1.021	2.395	3.661

Source: Field Study 2023

### **3.2 Result for Temperature Analyses**

The results of temperature data evaluation from 1981 to 2020 are presented in Table 2. The average annual temperature in Birniwa was 30.5°C, with a lowest of 29.3°C and a highest of 31.9°C. In Dutse, the temperature data indicated an average of 26.9°C, a lowest value of 25.5°C, and a highest value of 28.5°C. While in Kaugama's the mean temperature values is 25.2°C, maximum is 27.39°C, and minimum is 23.5°C. The average temperature value for Kirikasamma was 28.00C, with a minimum of 27.10C and a maximum of 29.10C. The temperature analysis results for Maigatari revealed the average yearly temperature as 27.9°C, with a lowest of 27.1°C and a highest

of 28.9°C. In Sule-Tankarkar's average yearly temperature was 27.7°C, with a lowest of 26.7°C and a highest of 28.9°C.

**Table 2. Mean Temperature Data**

Statistic	Birniwa	Dutse	Kaugama	Kirikasamma	Maigatari	Sule-Tankarkar
Nbr. of observations	40	40	40	40	40	40
Minimum	29.3	25.5	23.5	27.1	27.1	26.7
Maximum	31.9	28.5	27.4	29.1	28.9	28.9
Range	2.702	2.991	3.93	2.022	1.833	2.251
Mean	30.5	26.9	25.2	28.0	27.9	27.7
Variance (n-1)	0.596	0.764	1.343	0.315	0.242	0.358
Standard deviation	0.772	0.874	1.159	0.561	0.492	0.599
Variation coefficient	0.025	0.032	0.045	0.020	0.017	0.021
Skewness (Pearson)	0.286	0.280	0.481	0.297	0.064	0.210
Kurtosis (Pearson)	-1.030	-1.149	-0.918	-0.769	-0.710	-0.859

Source: Field Study 2023

### 3.3 Result for Wind Speed Analyses

The results for wind speed from 1981 to 2020 are illustrated in Table 5. Starting with Birniwa, the average wind speed was 1.4 m/s, while the minimum and maximum recorded values were 1.3 m/s and 1.5 m/s, independently. For Dutse, the average speed measured was 1.7 m/s, ranging from a minimum of 1.6 m/s to a highest of 1.8 m/s. The results further revealed that for Kaugama with average atmospheric speed of 1.5 m/s, with a minimum of 1.4 m/s and a maximum of 1.6 m/s. In Kirikasamma's the wind speed mean values was 2.4 m/s, with a lowest of 2.2 m/s and a highest of 2.6 m/s. The average annual wind speed at Maigatari was 3.5 m/s, with a lowest of 3.2 m/s and a highest of 3.7 m/s. Lastly, in Sule-Tankarkar an average yearly wind speed of 2.5 m/s, with a lowest of 2.2 m/s and a highest of 2.7 m/s was reported (Table 3).

**Table 3. Descriptive Statistics (Wind)**

<b>Statistic</b>	<b>Birniwa</b>	<b>Dutse</b>	<b>Kaugama</b>	<b>Kirikasamma</b>	<b>Maigatari</b>	<b>Sule_Tankarkar</b>
<b>Nbr. of observations</b>	40	40	40	40	40	40
<b>Minimum</b>	1.3	1.6	1.4	2.2	3.2	2.2
<b>Maximum</b>	1.5	1.8	1.6	2.6	3.7	2.7
<b>Range</b>	0.207	0.218	0.218	0.351	0.517	0.446
<b>Mean</b>	1.4	1.7	1.5	2.4	3.5	2.5
<b>Variance (n-1)</b>	0.003	0.003	0.003	0.008	0.020	0.010
<b>Standard deviation (n-1)</b>	0.052	0.056	0.055	0.089	0.140	0.100
<b>Variation coefficient</b>	0.037	0.033	0.035	0.037	0.040	0.040
<b>Skewness (Pearson)</b>	-0.291	0.019	-0.231	-0.134	-0.228	-0.165
<b>Kurtosis (Pearson)</b>	-0.614	-0.596	-0.722	-0.721	-0.864	-0.285

Source: Field Study 2023

### 3.4 Trend Analysis

The Mann-Kendall (MK) test (Mann 1945, Kendall 1975, Gilbert 1987) is utilized to numerically assess provided the outcome variable has shown repetitive growth or reduce gradually. A monotonic movement, regardless of its direction, signifies that the variable frequently grows (or lessens) gradually, even if the trend is not straightforward. The MK test is best viewed as a tool for exploratory analysis, according to Hirsch, Slack, and Smith (1982). It's most appropriate uses are spotting stations with significant or substantial changes and measuring these changes.

**Table 4. Mann-Kendall Trend Test Results**

<i>Location</i>	<i>Variables</i>	<b>Kendall's Tau (Z)</b>	<b>S</b>	<b>Var (S)</b>	<b>P-Value</b>	<b>Sens' Slope</b>	<b>Hypothesis</b>
<b>BIRNIWA</b>	<b>Rainfall</b>	-0.274	-	44.000	0.012	-7.432	H <sub>0</sub> = Rejected
	<b>Temperature</b>	0.538	420.000	0.000	0.0001	0.048	H <sub>0</sub> =Rejected
	<b>Wind</b>	-0.153	-	73.667	0.169	-0.001	H <sub>0</sub> =Accepted
<b>DUTSE</b>	<b>Rainfall</b>	-0.308	-	44.000	0.005	-11.842	H <sub>0</sub> =Rejected

	<b>Temperature</b>	0.494	385.000	7365.667	0.0001	0.052	H <sub>0</sub> =Rejected
	<b>Wind</b>	44.000	-86.000	44.000	0.324	-7.206E-4	H <sub>0</sub> =Accepted
<b>KAUGAMA</b>	<b>Rainfall</b>	-0.285	-	44.000	0.009	-9.135	H <sub>0</sub> = Rejected
			222.000				
	<b>Temperature</b>	-0.144	-	0.000	0.197	-0.039	H <sub>0</sub> =Accepted
			112.000				
	<b>Wind</b>	-0.183	-	7365.667	0.098	-0.001	H <sub>0</sub> =Accepted
			143.000				
<b>MAIGATARI</b>	<b>Rainfall</b>	-0.274	214.000	44.000	0.012	3.57	H <sub>0</sub> = Rejected
	<b>Temperature</b>	0.241	188.000	0.000	0.029	0.018	H <sub>0</sub> = Rejected
	<b>Wind</b>	-0.456	-	7365.667	0.0001	-0.008	H <sub>0</sub> =Rejected
			143.000				
<b>KIRIKASAMMA</b>	<b>Rainfall</b>	-0.062	-48.000	44.000	0.586	3.57	H <sub>0</sub> =Accepted
	<b>Temperature</b>	0.379	296.000	0.000	0.000	0.028	H <sub>0</sub> =Rejected
	<b>Wind</b>	-0.413	-	44.000	0.000	-0.005	H <sub>0</sub> =Rejected
			322.000				
<b>SULE-TANKARKAR</b>	<b>Rainfall</b>	-0.115	-90.000	44.000	0.302	-1.994	H <sub>0</sub> =Accepted
	<b>Temperature</b>	0.372	290.000	0.000	0.001	0.028	H <sub>0</sub> =Rejected
	<b>Wind</b>	-0.413	346.000	44.000	0.001	-0.005	H <sub>0</sub> =Rejected

**Source: Field Study 2023**

This study utilized time series analysis of three meteorological parameters (Rainfall, Temperature, and Wind) over a 39-year period (1981–2020) in seven LGAs of Jigawa state to identify the trend and slope magnitude of the series. A statistical test (Mann-Kendall test) (Table 4) and Sen’s Slope Estimator were utilized to analyze the trend. The results of Mann-Kendall’s Tau and the size of Sen’s Slope Estimator (see Fig. 2 a-f) indicated that the yearly trend analysis revealed both upward and downward trends for the variables across the LGAs.




Fig. 2a: Sen's Slope for Rainfall, Temperature, and Wind (Birniwa)




Fig. 2b: Sen's Slope for Rainfall, Temperature, and Wind (Dutse)




Fig. 2c: Sen's Slope for Rainfall, Temperature, and Wind (Kaugama)




Fig. 2d: Sen's Slope for Rainfall, Temperature, and Wind (Maigatari)




Fig. 2e: Sen's Slope for Rainfall, Temperature, and Wind (Kirikasamma)




Fig. 2f: Sen's Slope for Rainfall, Temperature, and Wind (Sule-Tankarkar)

The Mann-Kendall (MK) test results showed mixed outcomes regarding the null theory, which asserts that “there is no trend in the series” of the climatic variables (rainfall, temperature, and wind) being studied. For the rainfall series, with the exception of Kirikasamma and Sule-Tankarkar local government areas, the hypothesis was rejected. The null hypothesis for the temperature series was accepted solely for Kaugama local government area. On the other hand, in all local government areas except for Sule-Tankarkar, Kirikasamma, and Maigatari, the null hypothesis regarding wind speed was accepted. The rainfall trend is observed in only two out of seven local government areas, and the temperature trend is present in just one. This indicates that there are only variations, not changes, in the area's climate, as concluded from the synthesis of climate data over 39 years (1981–2020). Salam et al. (2021) corroborated this by indicating that from 1990 to 2020, the climate of Jigawa state exhibited significant earthly and geographical transfer in its deviation and alteration, which according to Akinsalona and Agunjobi (2014) is the changes in mean states and other statistical measures (like standard deviations, frequency of excessive, etc.) of the climate across all earthly and geographical scales yonder those of single climate phenomena (IPCC, 2018).

#### **4. Conclusion**

It reached the conclusion that the study area is challenged by irregularities in rainfall that are linked to heightened climate variability. Evidence was shown through the statistically significant spatial variations in rainfall, temperature, and wind speed, albeit at different scales. Overall, the findings exhibited a degree of spatial and temporal inclination for rainfall, temperature, and wind speed; therefore, the temporal variability is more evident than the spatial changes. This does not imply that spatial variations should be ignored, as they will have an impact on the livelihoods of farmers in the area. While precipitation portrays a significant part in climate change studies, temperatures

and wind are also significant as they affect evaporation and transpiration, which are key parameters of climate change.

## **5. References**

- Dan'azumi, S., & Ibrahim, U. A. (2023). Trend analysis of observed precipitation, temperature, and streamflow for Hadejia-Nguru wetlands catchment, Nigeria. *Theoretical and applied climatology*, *151*(1), 195-207.
- Ghalehtimouri, K. J., Ros, F. C., Rambat, S., & Nasr, T. (2024). Spatial and temporal water pattern change detection through the normalized difference water index (NDWI) for initial flood assessment: a case study of Kuala Lumpur 1990 and 2021. *Journal of Advanced Research in Fluid Mechanics and Thermal Sciences*, *114*(1), 178-187.
- Ishaku, D., Umaru, E. T., Adebayo, A. A., Löwner, R., & Okhimamhe, A. A. (2024). Analysis of the Observed Trends in Rainfall and Temperature Patterns in North-Eastern Nigeria. *Climate*, *12*(12), 219.
- Ledda, A., Serra, V., Ruiu, M. G. G., Bardi, A., Trogu, D., Di Cesare, E. A., & De Montis, A. (2024). Mainstreaming Climate Change Adaptation into Sectoral Plans: An Assessment Based on the Logical Framework Approach. *Sustainability*, *16*(9), 3705.
- Mustapha, A., Aris, A. Z., Yusoff, F. M., Zakaria, M. P., Ramli, M. F., Abdullah, A. M., . . . Narany, T. S. (2014). Statistical approach in determining the spatial changes of surface water quality at the upper course of Kano River, Nigeria. *Water Quality, Exposure and Health*, *6*(3), 127-142.
- Mwangi, H. M., Julich, S., Patil, S. D., McDonald, M. A., & Feger, K.-H. (2016). Relative contribution of land use change and climate variability on discharge of upper Mara River, Kenya. *Journal of Hydrology: Regional Studies*, *5*, 244-260.

- Ndabula, C., Averik, P., Jidauna, G., Abaje, I., Oyatayo, T., & Iguisi, E. (2013). Analysis of the spatio-temporal dynamics of landuse/landcover structures in the Kaduna innercore city region, Nigeria. *American Journal of Environmental Protection, 1*(4), 112-119.
- Njuguna, S. M., Onyango, J. A., Githaiga, K. B., Gituru, R. W., & Yan, X. (2020). Application of multivariate statistical analysis and water quality index in health risk assessment by domestic use of river water. Case study of Tana River in Kenya. *Process Safety and Environmental Protection, 133*, 149-158.
- O'Neill, B. C., Oppenheimer, M., Warren, R., Hallegatte, S., Kopp, R. E., Pörtner, H. O., . . . Licker, R. (2017). IPCC reasons for concern regarding climate change risks. *Nature Climate Change, 7*(1), 28.
- Pachauri, R. K., Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., . . . Dasgupta, P. (2014). *Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change: IPCC.*
- Pareek, A., Dhankher, O. P., & Foyer, C. H. (2020). Mitigating the impact of climate change on plant productivity and ecosystem sustainability (Vol. 71, pp. 451-456): Oxford University Press UK.
- Roy, J., Prakash, A., Some, S., Singh, C., Bezner Kerr, R., Caretta, M. A., . . . Tirado-von der Pahlen, M. C. (2022). Synergies and trade-offs between climate change adaptation options and gender equality: a review of the global literature. *Humanities and Social Sciences Communications, 9*(1).
- Sasanya, B. F., Adesogan, S. O., & Ademola, A. A. (2025). Spatio-temporal analysis of rainfall over Chad River Basin, Nigeria. *Theoretical and applied climatology, 156*(2), 118.
- Sawa, B., & Adebayo, A. (2011). The Impact of Climate Change on Precipitation Effectiveness Indices in Northern Nigeria.

- Tanko, A. I. (2013). 9 Agriculture, livelihoods and fadama restoration in northern Nigeria. *Wetland Management and Sustainable Livelihoods in Africa*, 205.
- Ujoh, F. (2014). Population Growth and Land Resource Conflicts in Tivland, Nigeria. *Resources and Environment*, 4(1), 67-78.
- Umar, A. T., & Bako, M. M. (2019). Recent Rainfall Trends and Variability in Sudano-Sahelian Region of Nigeria (1986-2015). *Ghana Journal of Geography*, 11(1), 33-57.
- Umar, D. A., Ramli, M. F., Tukur, A. I., Jamil, N. R., & Zaudi, M. A. (2021). Detection and prediction of land use change impact on the streamflow regime in Sahelian river basin, northwestern Nigeria. *H2Open Journal*, 4(1), 92-113.
- Umar, D., Umar, H., & Tukur, A. (2017). Climate variability and water supply: A review of rural water planning techniques for semi-arid region of Nigeria. *Dutse Journal of Pure and Applied Sciences (DUJOPAS)*, 3(2), 318-330.
- Umar, D. u. A., Garo, T. U., Lawan, M. M., Shehu, A. U., Dahiru, R., Adamu, A., & Babik, Z. M. (2025). Evaluating the effects of land use changes on stream-flow patterns in the semi-arid region of Nigeria. *African Research Reports*, 1(6), 537-546.
- Umar, D. A., Ramli, M. F., Aris, A. Z., Jamil, N., & Tukur, A. I. (2019). Surface water resources management along Hadejia River Basin, northwestern Nigeria. *H2Open Journal*, 2(1), 184-199.
- White, R., Uljee, I., & Engelen, G. (2012). Integrated modelling of population, employment and land-use change with a multiple activity-based variable grid cellular automaton. *International Journal of Geographical Information Science*, 26(7), 1251-1280.
- Zhao, J., Zhang, N., Liu, Z., Zhang, Q., & Shang, C. (2024). SWAT model applications: From hydrological processes to ecosystem services. *Science of the Total Environment*, 172605.

